


10 Warning Signs of Primary Immunodeficiency

Primary Immunodeficiency (PI) causes children and adults to have infections that come back frequently or are unusually hard to cure. 1:500 persons are affected by one of the known Primary Immunodeficiencies. **If you or someone you know is affected by two or more of the following Warning Signs, speak to a physician about the possible presence of an underlying Primary Immunodeficiency.**




Four or more new ear infections within one year.



Two or more serious sinus infections within one year.




Two or more months on antibiotics with little effect.



Two or more pneumonias within one year.




Failure of an infant to gain weight or grow normally.



Recurrent, deep skin or organ abscesses.



Persistent thrush in mouth or fungal infection on skin.



Need for intravenous antibiotics to clear infections.



Two or more deep-seated infections including septicemia.



A family history of PI.

Presented as a public service by:



Jeffrey Modell Foundation | Curing PI. Worldwide.



Funding was made possible in part by grant 5H75DP225146-05 from the United States Centers for Disease Control and Prevention (CDC)



National Heart, Lung, and Blood Institute (NHLBI)



National Institute of Allergy and Infectious Diseases (NIAID)



National Institute of Allergy and Infectious Diseases (NIAID)



National Institute of Child Health & Human Development

Baxter BioScience

CSL Behring Biotherapies for Life™

GRIFOLS

octapharma

PPTA Plasma Protein Therapeutics Association

Talecris BIOETHERAPEUTICS

These warning signs were developed by the Jeffrey Modell Foundation Medical Advisory Board. Consultation with Primary Immunodeficiency experts is strongly suggested. © 2010 Jeffrey Modell Foundation

For information or referrals, contact the Jeffrey Modell Foundation: 866-INFO-4-PI | info4pi.org