

## LOOKING FOR MORE OUT OF YOUR FACTOR IX (FIX) THERAPY?

Going in prepared helps make the most out of healthcare provider visits. Use this guide to kickstart the conversation.

- 1. What do you want your healthcare provider to know about your current treatment:** What's working? What's not?
- 2. Consider sharing how your treatment could better fit your lifestyle.** For example: "I want to know I'm protected from bleeds when I'm traveling or working out."
- 3. Ask your healthcare provider about keeping your trough levels higher for longer.**  
Factor IX trough levels affect many aspects of therapy, from bleed protection to dosing, so it's important to know yours and discuss them.
- 4. What about the future? Tell your healthcare provider about any upcoming life events and plans.**  
I would like my therapy to help me keep up with:
  - Achieving zero bleeds
  - Maintaining higher trough levels
  - Longer bleed protection
  - Dosing that works with my lifestyle
  - Treatment that matches my activities/goals
  - Other
- 5. What are the most important things you want to discuss with your healthcare provider?**

### **A point about joints**

People with trough FIX levels >15% have a very low risk for joint bleeds.

### **Trough levels: Every percent counts!**

According to the World Federation of Hemophilia, every added percentage point of baseline factor levels decreases bleeding frequency.

### **Notes:**

## IMPORTANT SAFETY INFORMATION

In clinical trials for IDELVION, headache and dizziness were the only side effects occurring in more than 1% of patients (1.8%), but are not the only side effects possible. Tell your healthcare provider about any side effect that bothers you or does not go away, or if bleeding is not controlled with IDELVION.

Please see full Important Safety Information on third page.



## ONLY IDELVION LETS YOU

**TAKE FIX LEVELS TO ANOTHER LEVEL**  
WITH 20% TROUGH LEVELS ON 7-DAY DOSING AND THE LONGEST SINGLE-DOSE HALF-LIFE OF ANY FIX†



**RECLAIM SPONTANEOUS**  
WITH 0 SPONTANEOUS BLEEDS‡



**FIT DOSING TO YOUR LIFESTYLE**  
WITH 7- AND 14-DAY DOSING OPTIONS§  
FDA-APPROVED

\*Hemophilia FIX Market Assessment. Third-Party Market Research.

†Based on a single dose of IDELVION at 50 IU/kg in adults (≥18 years).

‡The median AsBR for people who started and stayed on 7- or 14-day prophylaxis was 0. For people who switched to prophylaxis from on-demand, the median AsBR was 0.7. AsBR=annualized spontaneous bleed rate.

§Once well-controlled (1 month without spontaneous bleeding or requiring dose adjustments on a weekly dose of ≤40 IU/kg), people 12 years and older can be transitioned to 14-day dosing.



**Want to talk to someone who knows what you're going through?**

Connect with our Advocate Program<sup>||</sup>



**Call 1-866-705-9661**  
or **scan above**

<sup>||</sup>Advocates are not healthcare providers or medical experts and are compensated by CSL Behring for their time and expenses. For medical questions, you should contact your physician.

### Notes:

## IMPORTANT SAFETY INFORMATION

In clinical trials for IDELVION, headache and dizziness were the only side effects occurring in more than 1% of patients (1.8%), but are not the only side effects possible. Tell your healthcare provider about any side effect that bothers you or does not go away, or if bleeding is not controlled with IDELVION.

Please see full Important Safety Information on next page.



## IMPORTANT SAFETY INFORMATION

IDELVION®, Coagulation Factor IX (Recombinant), Albumin Fusion Protein (rFIX-FP), is used to control and prevent bleeding episodes in people with hemophilia B. Your doctor might also give you IDELVION before surgical procedures. Used regularly as prophylaxis, IDELVION can reduce the number of bleeding episodes.

IDELVION is administered by intravenous injection into the bloodstream, and can be self-administered or administered by a caregiver. Do not inject IDELVION without training and approval from your healthcare provider or hemophilia treatment center.

Tell your healthcare provider of any medical condition you might have, including allergies and pregnancy, as well as all medications you are taking. Do not use IDELVION if you know you are allergic to any of its ingredients, including hamster proteins. Tell your doctor if you previously had an allergic reaction to any FIX product.

Stop treatment and immediately contact your healthcare provider if you see signs of an allergic reaction, including a rash or hives, itching, tightness of chest or throat, difficulty breathing, lightheadedness, dizziness, nausea, or a decrease in blood pressure.

Your body can make antibodies, called inhibitors, against Factor IX, which could stop IDELVION from working properly. You might need to be tested for inhibitors from time to time. IDELVION might also increase the risk of abnormal blood clots in your body, especially if you have risk factors. Call your healthcare provider if you have chest pain, difficulty breathing, or leg tenderness or swelling.

In clinical trials for IDELVION, headache and dizziness were the only side effects occurring in more than 1% of patients (1.8%), but are not the only side effects possible. Tell your healthcare provider about any side effect that bothers you or does not go away, or if bleeding is not controlled with IDELVION.

**Please see full prescribing information for IDELVION, including patient product information.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

You can also report side effects to CSL Behring's Pharmacovigilance Department at 1-866-915-6958.

## CSL Behring

IDELVION is manufactured by CSL Behring GmbH and distributed by CSL Behring LLC. IDELVION® is a registered trademark of CSL Behring Lengnau AG.  
©2022 CSL Behring LLC 1020 First Avenue, PO Box 61501, King of Prussia, PA 19406-0901 USA